

## Study of Edaphic Factors of the Location for the Growth of White Sandal (*Santalum album L.*) in Indo-Nepal Border

<sup>1</sup>Roshan Kumar Yadav, <sup>2</sup>Subrata Mukhopadhyay and <sup>3</sup>Jagatpati Tah

<sup>1,3</sup>Department of Life Science and Biotechnology, Jadavpur University, 188 Raja Subodh Chandra Mullick Road, Jadavpur, Kolkata – 700 032, India.

<sup>2</sup>Department of Chemistry, Jadavpur University, 188 Raja Subodh Chandra Mullick Road, Jadavpur, Kolkata – 700 032, West Bengal, India.

Article Received: 20 April 2018

Article Accepted: 29 July 2018

Article Published: 24 August 2018

### ABSTRACT

Environment plays an important role for growing plants in any location in any country. White sandal (*Santalum album L.*) cultivation is also dependent on some specific edaphic factors along with host-parasite relationship etc. Soil environment factors play a vital role for the growth of white sandal was unknown to the farmer's community though it grows in wild condition since ancient civilization. It is common belief to people that white sandal plant cannot survive without proper environment and specific host plants surrounding its root zone. Keeping all these views in mind an experiment was undertaken in some areas (Rampur, Malangwa, Kamalpur, Bardibas) in Indo-Nepal Border. Soil samples were taken from each and every forest garden and analysed in the laboratory to assess the soil nutrient status. The aims and objects of the study were to observe the role of soil nutrients of the locations in favour of the growth and development of *Santalum album L.*

**Key words:** Edaphic factors, Environmental factors, Root zone, Nutrient status.

### INTRODUCTION

Sandalwood (White Sandal) is the fragrant heartwood of some species of genus *Santalum*. The widely distributed and economically important *Santalum* genus belongs to the family Santalaceae which includes 30 genera with about 400 species, many of which being completely or partially parasitic (John, 1947). The word Sandal has been derived from Chandana (Sanskrit), Chandan (Persian), Savtador (Greek) and Santal (French). There are references of Sandalwood in Indian mythology, folklore and ancient scripts. 'Chandana' the Sanskrit name ascribed to *Santalum album L.* was known and used in India from the earliest historic times and is frequently mentioned in the ancient Sanskrit writings, some of which dated before Christian era. Kautilya's Arthashastra (320 B.C.) considered Sandal as one of the important forest products to increase royal revenue. Charaka Sanhitha, the major text book of internal medicine in Ayurveda (300 B.C.) quotes uses of Sandal over 160 time in the entire text. In treatment of major diseases like fever, piles, hemorrhagic conditions, diabetes, dropsy, mental disorders, management of poisons & skin disorders wide spread uses of sandal is seen. Susrutha Samhita (150 B.C.) a great text on Indian wisdom on surgical procedures, equally preferred sandal for the management of wounds. Sandal fumigation is indicated in warding off evils and organisms, which contaminate the wounds. Such fumigations hasten the wound healing & surgical wards remain aseptic. Dusting of wounds with sandal for early healing is common. In the Amarkosha (Lexicon 3<sup>rd</sup> or 4<sup>th</sup> Century A.D.) sandal is mentioned and it is said that 'Vina-malayam anyathra chandanam vivarditha' [Majumdar, 1941].

The extraction and disposal of sandal came under the Forest Department in 1864 in Mysore state [Adkoli, 1977]. In Karnataka (formerly Mysore) the forest working plan for sandal extraction were prepared for Hunsur Talik in 1910, Heggadadevanakote in 1920 and Narasimharajapura in 1926. In 1871, the parasitic nature of sandal was reported by John Scott. Watt (1893) described the technique of raising sandal seedlings in tile pots in the nurseries and planting in the field. McCarthy (1899) first noticed the spike disease of sandal in Coorg. Brandis (1903) suggested

that though sandal is a root parasite, it may derive part of its nutrition from the soil as well. Barber (1905) noted that haustoria formation occurred only on certain roots of sandal and not on all of them.

Ecologically sandal has adapted various agro - climatic and soil conditions for *in situ* regeneration with an exception of waterlogged areas and very cold places. In India, 8 Sandal growing areas have been identified as potential provenances of Sandal on the basis of population density, phenotypic characteristics, latitude, longitude and eco-climate (Jain *et al.*, 1998). The provenances vary in climate and edaphic preference since they are located in different localities of South and Central India. The state of West Bengal is cited in the map of occurrence and distribution of *Santalum album* in India (Srinivasan *et al.*, 1992).

Some hindrances and problems for its propagation were observed by various workers in these areas. Keeping all these views in mind we are going to undertake the venture for its mass propagation through seeds and its cultivation with their agronomical maintenance properly.

## MATERIALS AND METHODS

### Materials

1. Sandalwood seeds: Seeds of *Santalum album* L. were collected from Hirbunndh mouza of Hirbunndh Range under Bankura ( South ) Forest Division during the month of November-December and May-June of 2015 and 2016 for experimentation. Simultaneously, seeds of *Santalum album* L. were also procured from Institute of Wood Science & Technology, Bangalore in the month of February, 2015 for the same experimentation purposes.
2. Chemicals: Gibberelic acid (GA<sub>3</sub>)
3. Apparatus: Container, Markin cloth, Polypots, Hycopots.
4. Miscellaneous: Sand, bricks, Seive, FYM, Water, etc.
5. Meteorological Informations

**Table I :** Temperature regime of the experiment site (in degree Celsius)

Months	2015		2016		2017	
	Max	Min	Max	Min	Max	Min
JAN	27	09	27	9	27	9
FEB	32	11	34	9	32	9
Mar	39	14	38	15	37	16
APR	43	20	42	9	38	18
MAY	44	24	45	22	43	21
JUN	39	24	37	22	41	23
JUL	38	24	37	25	35	25
AUG	35	23	34	24	35	25

SEP	34	24	36	24	35	22
OCT	34	16	35	20	34	18
NOV	32	11	32	13	32	16
DEC	29	10	29	7	27	9
IN YEAR	44	9	45	7	43	9

**Table II :** Monthly rainfall in the district of the experiment site (in degree Celsius)

Month	Normal			
		2015	2016	2017
January	17	20	3	7
February	12	13	40	15
March	19	7	9	4
April	27	4	57	66
May	65	52	76	77
June	198	213	247	183
July	272	331	255	471
August	293	332	445	386
September	246	358	139	384
October	122	18	95	0
November	15	99	00	0
December	03	6	00	0
Total for the year	1289	1453	1366	1687

#### 6. Information on Sites

Abbreviation	Full name	Total
L	Location	4
P	Plantation	3
T	Treatment	9
Y	Year	3

#### 7. Name of Location

No. of Location	Name of Location
L <sub>1</sub>	Rampur

L	Malangwa
L <sub>3</sub>	Kamalpur
L <sub>4</sub>	Bardibas

#### 8. Information of Treatments

No. of Treatment	Composition [Host(S)]
T <sub>0</sub>	No Host
T <sub>1</sub>	Arhar
T <sub>2</sub>	Tulsi
T <sub>3</sub>	Arhar+Tulsi
T <sub>4</sub>	Akand
T <sub>5</sub>	Arhar+Akand
T <sub>6</sub>	Ghantu
T <sub>7</sub>	Arhar+Ghantu
T <sub>8</sub>	Nayantara
T <sub>9</sub>	Tulsi+Nayantara

#### 9. Information of Plantation

No. of Plantation	Year
P <sub>1</sub>	2015
P <sub>2</sub>	2016
P <sub>3</sub>	2017

10. Growth Parameters, viz. plant height (cm) and basal girth (cm), were measured out of all planted saplings year-wise from each and every forest garden. *Santalum album* L.

## METHODS

### A) Germination Study in Nursery:

**i) Pretreatment by soaking in water:** Sandalwood seeds are soaked in water for 24 hours before sowing. Seeds are sown in sand bed (6 mm deep). Germination starts after 60 days. In-between 61 to 100 days, only 3-4% germination is obtained.

**ii) Pretreatment by boiling water:** Sandal seeds are pretreated with boiling water (10 parts of boiling water with one part of seeds) for 1 minute and then kept in normal water overnight for soaking. Treated seeds are sown in sand bed. Germination starts after 50 days. In-between 51 to 100 days, hardly 8% germination is obtained.

**iii) Pretreatment by alternate wetting & drying:** Sandal seeds are exposed to alternate wetting & drying for 12 hours wetting followed by 12 hours drying in sun. This process is repeated for 7 days and then the seeds are sown

in sand bed (6 mm deep). Germination starts after 40 days. In-between 41 to 100 days, 8 to 9% germination is obtained. Few germination is found even after 100 days upto 150 days.

**iv) Pretreatment by Gibberellic Acid:** Matured seeds were collected from the sandalwood trees of Hirbandh Block in November- December, depulped, dried in sun and stored in polybags for germination test. 6 samples each of 300 sandalwood seeds were taken and tied in markin cloth. 3 containers (1 liter each) were taken for 3 different concentrations (0.0125%, 0.025%, 0.05%) of gibberellic acid. 2 seed samples were dipped in each container for 16 hours & 24 hours soaking. The treated sandalwood seeds were then sown in the sand bed. The sand beds were watered twice daily in the morning & afternoon. First germination was started after 24 days of seed sowing. The number of seeds germinated in each treatment is recorded and the germination is continued upto 90 days after sowing. The germinated seedlings were transplanted in polypot (8"x4" & 9"x 5") and 300 cc hycopots in nursery at 3 to 4 leaf stage.

Sandal seeds have a post drop dormancy of two months due to presence of hard seed coat or due to presence of chemical substances in the seed coat which are impervious to water and gases. Germination of Sandal seeds are found profuse from the bird droppings in the forest floor as well as in the village yards and bunds of the agricultural fields. Sandal is also found growing wild in some farmlands, homesteads and wastelands in Hirbundh block of Bankura District and Arabari forests of Midnapur District. This indicates the potential of growing the tree in the farm lands.

However, germination of seeds are very low (10-15% within 60 days) when the seeds are sown in mother bed (sand beds) after hot and cold water treatment or alternate wetting & drying due to its hard seed coat and dormancy. Sandal seeds have been found to germinate faster when the seed coat is completely removed, or seeds are soaked in 0.05% gibberellic acid for 12-16 hours (Nagaveni and Srimathi, 1981). In sandal seeds, the duration of germination is much prolonged after the dormancy period. It starts in 25 days and reaches hardly 50% in 90 days with 0.05% GA<sub>3</sub> soaking for 16 hours (Das and Tah, 2013). Germination study conducted in Hirbundh & Kamalpur nurseries of Bankura (South) Division in 2011, 2012 & 2013 where seeds are soaked in different concentration (0.0125%, 0.025%, 0.05%) of gibberellic acid for 16 hours & 24 hours respectively and sown in sand beds of nursery. The germinated seedlings are pricked out from sand bed and planted in polypot & hycopot beds of nursery at 3 to 4 leaf stage and kept in nursery without & with host (*Cajanus cajan*).

Plantation of Sandal plant in different forest gardens in South West Bengal

Sl. No.	Location	Year of Planting	No. of seedlings Planted
1	Rampur	2015	100
2	Malangwa	2015+2016	150+ 200
3	Kamalpur	2016	30
4	Bardibas	2016	100

## **B) Soil Testing Methods:**

### **(i) Soil pH**

The pH of the soil was determined with the help of a pH meter in 1: 2.5 soil: water suspension ratio as described by Jackson (1973).

### **(ii) Oxidizable Organic Carbon**

Organic carbon was determined by oxidizing soil with 1 (N) potassium dichromate  $K_2Cr_2O_7$  in presence of concentrated  $H_2SO_4$  and back titrating the remaining  $K_2Cr_2O_7$  with ferrous ammonium sulphate solution using diphenylamine indicator, following the wet digestion method of Walkley and Black as described by Jackson (1973).

### **(iii) Available nitrogen (N)**

The available nitrogen (N) of the soil was estimated through the hot alkaline potassium permanganate method as suggested by Subbiah and Asija (1956).

### **(iv) Available phosphorus (P)**

Available phosphorus (P) of soil is determined by using Olsen's method. In this method, the extractant is 0.5M  $NaHCO_3$  solution adjusted to pH 8.5 with 10% NaOH.

### **(v) Available potassium (K)**

Available potassium content of the soil was determined by flame photometer after extraction with neutral normal ammonium acetate solution as described by Jackson (1973).

### **(vi) Available Zinc(Zn), Copper(Cu), Manganese(Mn) & Iron(Fe)**

The method developed by Lindsay and Norvell (1978) using DTPA (Diethylene Triamine Penta Acetic Acid) for separating soils into deficient and non-deficient categories for Zn, Cu, Mn & Fe is useful and adopted in this Laboratory analysis.

## **Extracting Solution**

To prepare 1 litre of DTPA extracting solution, dissolve 13.1 ml reagent grade TEA, 1.967 g DTPA (AR grade) and 1.47 g of  $CaCl_2$  in 100 ml of glass distilled water. Allow sometime for the DTPA to dissolve and dilute to approximately 900 ml. Adjust the pH to  $7.3 \pm .05$  with 1:1 HCl while stirring and dilute to 1 litre. Addition of approximately 4 ml of 1N HCl will bring the pH of the solution to 7.3. This solution is stable for several months.

## **Extraction and Determination**

Weigh 10 grams of air dried soil in a 125 ml conical flask or polypropylene bottle. Then add 20 ml of the DTPA extracting solution. Cork the bottles or flask and place them upright on a horizontal shaker. Shake for two hours with a speed of 120 cycles per minute. Filter the suspension through Whatman no 42 filter paper. Keep the filtrate in polypropylene bottles to be analysed for Zn, Cu, Mn and Fe with an atomic absorption spectrophotometer. Analyse

the sample as described above under standard curve. When samples need dilution before measurement, they should be diluted with DTPA solution to maintain a constant matrix. Experimental condition such as shaking time, DTPA concentration, pH and temperature during shaking influence the amount of Zn, Cu, Mn and Fe extracted by DTPA. The most suitable pH of extracting solution is 7.3, shaking time 2 hours and temperature during shaking  $25 \pm 1^{\circ}\text{C}$ . The values of the nutrient extracted will change if these precautions are not followed. Increase in shaking time and temperature markedly enhance the extractability of Zn, Cu, Mn and Fe. The increase in pH from neutral (7.0) to 7.9 had no effect on Zn and Cu while extractability of Fe and Mn was increased markedly.

**Zinc Standard Solution :** Dissolved 0.439 g AR grade Zinc sulphate  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$  in 200ml of glass distilled water in a beaker. Added 5 ml of 1:5  $\text{H}_2\text{SO}_4$ . Transfer to a litre measuring flask and made volume to the mark to have a standard solution of 100 ug Zn/ml (100 ppm). Transfer 10 ml of this standard solution to 100 ml volumetric flask and diluted to the mark with DTPA extracting solution to have a stock solution of 10 ug Zn/ml (10ppm). For preparing working standard, transfer 1,2,4 and 6 ml of stock solution(10 ug Zn/ml) to a series of clean 100 ml volumetric flask and each to the mark with DTPA extracting solution.

Volume of stock Zn solution taken : 0 1 2 4 6ml

Concentration of Zn now in solution: 0 0.1 0.2 0.4 0.6ug/ml (ppm)

**Iron Standard Solution :** Dissolved 0.702g of AR grade ammonium ferrous sulphate  $(\text{NH}_4)_2\text{SO}_4 \cdot \text{FeSO}_4 \cdot 6\text{H}_2\text{O}$  in 300 ml deionized or glass distilled water in a beaker. Added 5 ml of 1:5  $\text{H}_2\text{SO}_4$ . Transfer to a litre measuring flask and made volume to the mark. This is a standard solution of 100 ug Fe/ml (100 ppm). To prepare working standards, transfer 1,2,4 and 6 ml of stock solution and diluted each to the mark with DTPA extracting solution.

Volume of stock Fe solution taken : 0 1 2 4 6ml

Concentration of Fe in solution : 0 1 2 4 6ug/ml (ppm)

**Manganese Standard Solution :** Dissolved 0.288 g potassium permanganate ( $\text{KMnO}_4$ ) AR grade in 300 ml deionized water in a breaker. Added 20 ml concentrated  $\text{H}_2\text{SO}_4$  warm to about  $60^{\circ}\text{C}$  and added oxalic acid solution dropwise to make the solution colour less. Cool and transfer to a litre measuring flask and made volume to the mark. This solution contains 100 ug Mn/ml (100 ppm). To prepare working standards, transfer 1,2,4,6 and 8 ml of the standard solution to a series of clean 100 ml volumetric flask and diluted each to the mark with DTPA extracting solution.

Volume of stock Mn solution taken : 0 1 2 4 8 ml

Concentration of Mn in solution : 0 1 2 4 8ug/ml (ppm)

**Copper Standard Solution:** Dissolved 0.392 Copper sulphate ( $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ) of AR grade in 400 ml glass distilled water in a beaker. Transfer to a litre measuring flask and made volume to the mark with glass distilled water. This is a standard solution containing 100 ug Cu/ml. To prepare the working standards, transfer 1,2,4 and 6



ml of stock solution to a series of clean 100 ml volumetric flask and diluted each to the mark with DTPA extracting solution.

Volume of stock Cu solution taken : 0 1 2 4 6 ml

Concentration of Cu in solution : 0 1 2 4 6ug/ml (ppm)

#### **(vii) Analysis for Available Molybdenum (Mo)**

Ammonium oxalate (pH 3.3) or Grigg's reagent is considered to be the best one for the determination of available Mo. Further, this extractant is easy to prepare, have sufficient buffering capacity to prevent any material change in the pH of the soil extract and forms stable complexes with Mo ( $\text{MoO}_3 \cdot \text{C}_2\text{H}_2\text{O}_4$  and  $(\text{MoO}_3)_2 \cdot \text{C}_2\text{H}_2\text{O}_4$ ) Molybdate absorbed on soil colloids and clay is presumably replaced by the oxalations. This exchange is made irreversible by the formation of strong Mo-oxalic acid complexes to make a single extraction effective.

#### **Extracting Solution**

To prepare ammonium oxalate (pH 3.3), dissolve 24.9 g AR ammonium oxalate and 12.6 g oxalic acid per litre of solution, adjusted to pH 3.3.

#### **Standard Solution**

Prepare a solution containing 100 ug Mo/ml by dissolving 0.150 g AR grade  $\text{MoO}_3$  in 100 ml of 0.1N NaOH solution rendering it acidic with HCl and make the volume to 1 litre. Take 10 ml of this stock solution in one litre volumetric flask and make the volume to the mark to get a working standard of 1 ug Mo/ml. To make working standards, transfer 0.5,1.0,2.0,3.0 and 4.0 ml of 1 ug Mo/ml solution to series of clean 125 ml separatory funnels.

#### **Extraction and Determination of Molybdenum**

Twenty five gram soil is taken in a 500 ml corning or pyrex glass conical flask. To this 250 ml ammonium oxalate of pH 3.3 is added. The contents of the flask are shaken over an end to end horizontal shaker for 10 hours and filtered through Whatman filter paper no 50. Then, 200 ml of the filtrate is taken in a 250 ml breaker and evaporated to dryness on a water bath. The contents of the breaker are headed at high temperature ( $500^\circ\text{C}$ ) in a furnace for 5 hours to destroy organic matter and oxalates.

Contents of the breaker are then digested with 5 ml of diacid ( $\text{HNO}_3 - \text{HClO}_4$  4:1) after keeping it overnight, then with 10 ml of 4N  $\text{H}_2\text{SO}_4$  and  $\text{H}_2\text{O}_2$  every time taking to dryness. Then 10 ml of 0.1N HCl is added to the breaker and filter paper washed with another 10 ml. The filtrate is stored in corked plastic reagent bottles for estimation of available Mo by the procedure described above under standard curve.

#### **(viii) Analysis for Available Boron (B)**

Out of several methods devised to assess the level of available B in soil, the hot water soluble B method of Berger and Troug (1939) has been most widely accepted. Recent description of this method include some changes (Keren



and Bingham 1985, Sippola and Ervco 1987), but the basic procedure remains the same. Throughout boron analysis, use of borosilicate glassware should be avoided even for storage of chemicals. Plastic containers or corning pyrex glassware should be used.

### Standard Solution

To prepare the standard stock solution, dissolve 0.570 g boric acid ( $H_3BO_3$ ) AR grade in a litre distilled water to obtain a stock solution of 100 ug B/ml. Take 5 ml of the stock solution in a 100 ml volumetric flask and dilute to the mark. This solution contains 5 ug B/ml.

### Extraction and Determination of Boron

A 25 gm soil sample, 50 ml of water and about 0.5 g of activated charcoal is boiled for 5 minutes in a quartz flask and filtered immediately through Whatman filter paper No. 42. 5 ml of the extract is taken in a 25 ml volumetric flask and 4 ml of buffer masking solution and 4 ml of azomethine -H reagent solution is added. The colour is allowed to develop for 1 hour, and the volume is made to the mark. Intensity of colour is measured spectrophotometrically at 420 nm, and the B concentration read off from standard curve described above.

### E) Statistical Models:

Statistical Models and methods were done as followed by Singh & Chaudhary (1995) and Panse & Sukhatme (2005).

## RESULTS AND DISCUSSION

### Results

#### Soil Parameters Study

Six soil samples were collected from the different forest gardens of Bankura and Burdwan district, viz. Bagaldhara, Rangamati, Kamalpur, Beliatore & Hirbandh of Bankura and Khandari of Burdwan. The soil samples were tested for pH, Organic carbon, available N, P and K and presented in Table-III. The result show that Bagaldhara has highest macronutrient content and lowest is Kamalpur.

Table - III : Soil Test Reports of Macronutrients

Location	pH	OC (%)	Available N ( $Kg\ ha^{-1}$ )	Available N (PPM)	Available K (PPM)	Available P (PPM)
Rampur	5.71	0.51	234.6	106.63	57.6	8.2
Malangwa	5.70	0.47	221.3	100.59	51.8	7.9
Kamalpur	5.57	0.34	198.8	90.36	49.8	5.7
Bardibas	4.95	0.35	211.3	105.51	50.5	5.6
Rampur	5.66	0.44	217.7	98.95	51.7	7.8
Malangwa	5.30	0.42	210.8	95.82	52.1	5.5

Micronutrient analysis was done for 4 soil samples, viz. Bagaldhara, Rangamati, Kamalpur and Hirbandh and reflected in Table-XV. It is evident from the results that the micronutrient (Cu,Zn,Mn,Fe,Mo & B) content is lowest in Bagaldhara, Cu content is maximum in Rangamati, Zn & Fe content is highest in Kamalpur, Mn content is highest in Hirbandh. In all the cases B content is below the critical level (0.3 ppm) and Mo content is below detection level (BDL).

Table - IV : Soil Test Report of Micronutrients

Sl.No	Test Parameters	Critical levels of micronutrients (mgkg <sup>-1</sup> or PPM)	Results			
			Hirbandh	Kamalpur	Rangamati	Bagaldhara
1	Copper (Cu)	0.2	0.67	1.08	1.38	0.50
2	Zinc (Zn)	0.6	0.32	0.86	0.46	0.19
3	Manganese (Mn)	2.0	73.72	45.89	69.50	25.89
4	Iron (Fe)	4.5	8.66	17.15	7.96	2.25
5	Molybdenum (Mo)	0.05	BDL	BDL	BDL	BDL
6	Boron (B)	0.3	0.17	0.14	0.17	0.14

## DISCUSSION

### Study of soil parameters:

Sandal requires good drainage and does not stand waterlogging. Best growth of sandal trees are found on rich fairly moist soil such as garden loam and well drained deep alluvium on the river banks (Troup,1921). A significant relationship between available nitrogen content in 'A' horizon and annual growth increment was observed in the soils of Talamalai Range (Krishnamurthy *et al.*,1983). In a study carried out by Jain *et al.*,(1988) on soil properties and their relationship to the growth of sandal in three areas, it was observed that lime status, water holding capacity, pore space, volume expansion on wetting, exchangeable calcium and magnesium and available potash, exert positive influence on the increment in girth and height. Requirement of host for proper growth of sandal was demonstrated in a field study by Ananthapadmanabha *et al.*, (1984). Further analysis of soil and leaf samples from this trial had shown that sandal depends on its host for K, P and Mg (Rangaswamy *et al.*, 1986b). Sandal can draw other nutrients directly from soil because its roots have good cation exchange capacity (Parthasarathi *et al.*, 1971).

## REFERENCES

- Adkoli, N. S. [1977]. Sandalwood in Karnataka. *Retrospect and prospect – Proceedings of the All India Sandal Seminar*, Bangalore, 86-94.
- Ananthapadmanabha H.S., C.R.Rangaswamy, C.R. Sarma, H.C.Nagaveni, S.H. Jain, K.R. Venkatesan and H.P. Krishnappa (1984). Host requirement of sandal (*Santalum album* L.). *Indian Forester* 110(3) : 264 – 68.
- Barber, C.A.[1903]. Report on spike disease of sandalwood trees in Coorg. *Indian Forester*, 29: 21-31. Barber, C.A.[1903]. Studies in root parasitism – the houstorium of *Santalum album* L. 1. Early stages upto penetration. Mem. Dept. Agri,Ind. Bot.Soc.I., Pt. 1:1-26.



- Barber, C.A.[1902]. The natural history of sandal trees. *Ind.For.* 28: 340-41
- Barber, C.A.[1903] The mature houstorium. *Ibid J. Pt. 4* : 1-49.
- Barber, C.A.[1905]. The haustoria of sandal roots. *Indian Forester* 31: 189.
- Barber, C.A.[1906]. Studies in root parasitism: The haustoria of *Santalum album* L. *Mem Dept. Agri. Ind Bot. Ser 1* Pt. 1: 1-26
- Barber, C.A.[1907]. Studies of root parasitism: the haustoria of *Santalum album* L. *Mem Dept. Agri. Ind Bot. Ser 1* Pt., 11: 1-58.
- Berger, K.C. and E.Troug (1939). Boron determination in soils and plants. *Ind. Eng. Chem. Anal. Ed.* 11: 540-545
- Brandis, D. [1903]. Treatment of the sandal tree. *Indian Forester*, 29: 3-6.
- Das, S.C. and J. Tah (2013). Effect of GA<sub>3</sub> on seed germination of sandal (*Santalum album* L.). *International Journal of Current Science* 2013, 8 : E 79-84.
- Jackson, M. L. (1973). *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd.; New Delhi. Pp. 498-.
- Jain, S.H., V.G. Angadi, A.N. Rajeevalochan, K.H. Shankaranarayana, K.S Theagarajan and C.S. Rangaswamy.(1998). *Identification of provenances of Sandal in India* for genetic Conservation. In: *Sandal and its prodncts*, Raomiljac A.M., H.S.A padmanabha, R.M. Welbourn and K.S. Rao (eds.), Proceedings of an International Seminar held at bangalore, India. 18-19 December, 1997. Canberra, ACIAR Proceedings, 84 :117 – 120.
- John, H. (1947). The history, present distribution and abundance of sandalwood on Oahu, Hawaiian Islands. *Hawaiian Plant Studies Pacific Science. I:* 5- 20.
- Keren, R. and F.T. Bingham (1985). Boron in water, soils and plants. *Adv. Soil Sci.* 1: 229-276.
- Krishnamurthy, R., S. Kondas, A. Sekaranandam and P. Krishnamurthy. (1983). A Study on the performance of Sandal in the soils of Talamalai Range. *Indian J. of Forestry*, 6 : 17-23.
- Majumdar, G. P. [1941]. The white sandal. *Science and Culture*, 6: 492-495.
- McCarthy, C. [1899]. Progress Report of Forest Administration in Coorg for 1898-1899.
- Nagaveni, H.C & R.A. Srimathi. (1981). Sandal seeds, viability, germination and storage. Second All India Sandal Seminar, Salem. Tamil Nadu Forest Dept. 9-10, May.
- Nagaveni, H.C & R.A. Srimathi. (1981a). Studies on germination of the Sandal *Santalum album* L. Pretreatment of sandal seeds. *Indian Forester*, 107 (6) : 348 – 354.
- Panse, V.G. and P.V. Sukhatme (1995). *Statistical Methods for Agricultural Workers*. Published by I.C.A.R., Anusandhan Bhavan, Pusa, New Delhi,110012.
- Parthasarathi, K., S. K. Gupta and P. S. Rao [1971]. Studies on sandal spike, Part –IX Cation exchange Capacity of Sandal (*Santalum album* L.)in health and disease. *Curr. Sci.* 23 : 640 - 641.
- Rangaswamy, C.R., H.S.Ananthapadmanabha, S.H. Jain and H.C. Nagaveni (1986b). Nutrient uptake and host – requirement of sandal. *Van Vigyan*, 24: 75 -79.
- Singh, R.K.and Chaudhary,B.D.(1995) *Biometrical Methods in Quantitative Genetic Analysis*, Kalyani Publishers, New Delhi, 110002.

Srinivasan, V.V., V. R. Sivaramakrishnan, C. R. Rangaswamy, H.S. Ananthapadmanabha and K.H.[1992]. Sandal – *Santalum album* Linn. Indian Council of Forestry Research and Education. Publsd. ICFRE, Bangalore.

Subbiah, B.V. and Asija, G.L.(1956). A rapid procedure for the estimation of available nitrogen in soils. Current Science, 25: 259-260.

Watt, J. S. [1893]. *Economic products of India*.

Table-V: Some field photographs

<p><b>Plate-1: Sterilization of seeds before imbibe</b></p>	<p><b>Plate-2: imbibed seeds</b></p>
<p><b>Plate-3: Collection of data</b></p>	 <p><b>Plate-4: Seed sowing in the sand bed</b></p>
 <p><b>Plate-5: One year grown plant</b></p>	 <p><b>Plate-6: Seedling growing in hycopot</b></p>