

ARTHRAID Tablet –Advanced Nutritional Arthritis Formula

Govind Shukla, Anusha Palkamshetti, K.Srinidhi & C.J. Sampath Kumar

Lactonova Nutripharm (P) Ltd, Makers of ARTHRAID Tablets, 81/3, IDA Mallapur, Hyderabad, Telangana, India-500 076.

Article Received: 19 August 2018

Article Accepted: 18 November 2018

Article Published: 09 January 2019

ABSTRACT

Osteoarthritis is a chronic, inflammatory joint disease in the world. In India more than 20% of total population is suffering from arthritis, although the main cause of disease is unknown, morphological changes observed in OA include cartilage erosion as well as inflammation. Complex network of risk factors and biochemical parameters, including cytokines, proteolytic enzymes trigger the disease, by knowing the exact mechanism of progressive of disease, it may help in finding the new drug for reducing pain and curing of the joint disease. Conventional medicines do not prevent progression of osteoarthritis. Natural supplements have gained importance as a better alternative for the prevention of osteoarthritis progression. Eating a Natural supplement like Arthraid tablet can provide all the necessary nutrients and helps prevent progression of osteoarthritis. The present paper Reviews the Role of Arthraid tablets developed by R&D cell of Lactonova Nutripharm Pvt Ltd. Hyderabad in prevention of osteoarthritis progression.

Keywords: OA, Morphological changes, Cytokines, Arthraid tablet.

1. INTRODUCTION

Osteoarthritis (OA) is degenerative joint disease, which affects millions of people in the world. It is a complex disease whose pathogenesis, changes the tissue homeostasis of articular cartilage and subchondral bone, determine the predominance of destructive processes. A key role in the pathophysiology of articular cartilage is played by cell/extra-cellular matrix (ECM) interactions.




Fig.1

2. SIGNS AND SYMPTOMS

Findings from studies indicate that age, gender, joint impairment, reduced range of motion (ROM), joint stiffness, and pain, contribute to increased disability.1, 2

2.1. Pain

The most common symptom is a chronic pain, during development of knee joint inflammation the concentration of Excitatory amino acids (EAA) especially Glutamate is increased which is released from sensory neurons in the spinal cord contribute to hyperalgesia and pain in the affected area. Several studies have found that there is no correlation between radiological images and pain parameters, but the medial side of the knee showed most sensitization in patients with strong/severe knee OA, the degree of pain can be measured with temporal summation of pressure pain instrument.³⁻⁵

2.2. Joint stiffness

The concept of joint stiffness in arthritis and related pathology diseases was introduced in the early 1960s.^{6, 7} It is revealed that surface-active phospholipid (SAPL) (synovial surfactant) capable of reducing friction to the very low levels and provide lubricant in normal joint moreover, this lining is deficient in osteoarthritis and lead to stiffness of joint.^{8, 9}

2.3. Muscle weakness

Quadriceps muscle strengthening is an important protective function at knee joints. Cross-sectional studies suggest that strength is correlate with physical function and that increasing quadriceps strength reduces pain and improves function. Evidence suggests that thigh muscle strength may protect against knee joint damage and progression of existing OA.^{10, 11} Arthrogenic muscle inhibition (AMI) is a presynaptic, constant reflex inhibition of musculature surrounding a joint after damage to joint as it restricts full muscle activity and prevent the quadriceps strengthening, weaker quadriceps have been associated with an increased rate of loading at the knee joint.¹² AMI is caused by activity in multiple inhibitory pathways, its severity may vary according to the degree of joint damage.¹³

2.4. Bone enlargement and swelling

Due to pathological changes of articular cartilage in knee joint resulted from many causes leads to blockage and edema of soft tissues, disturbance of blood circulation, erosion and injury of chondrocyte, and even increase of bony density and formation of cystic changes, resulting in swelling and pain.¹⁴

3. RISK FACTORS OF KNEE OSTEOARTHRITIS

OA has a multifactorial etiology, can be considered the product of interaction between systemic and local factors.

4. SYSTEMIC RISK FACTORS FOR OA

4.1. Age

It is most important factor for development of osteoarthritis; with increasing age the tensile property of cartilage in articular cartilage is decreased results in accumulation of glycation which causes mechanical failure.¹⁵

4.2. Gender

Women have a higher level of pain and disability than men.¹⁶ A hospital-based study revealed rates of osteoarthritis is as high as 68% in women and 58% of men aged 65 and older.¹⁷

4.3. Genetics hormones

Classic study of monozygotic (MZ) twins aged 48 to 70 years, having identical genes showed 65% influence of genetic factors in developing of osteoarthritis.¹⁸ Between 39% and 65% of osteoarthritis in the general population can be attributed to genetic factors, women after menopause are more susceptible to knee arthritis because of increasing level of osteocalcin and bone resorption.¹⁹ Levels of osteocalcin, a marker of bone turnover, were lower in women with knee osteoarthritis.²⁰

4.4. Diet

Rapid changes in diet and lifestyle by consumption of unrefined carbohydrates and Junk foods increased the rate of chronic diseases.²¹ Furthermore, chondrocytes are powerful sources of reactive oxygen species, which may damage cartilage collagen and synovial fluid hyaluronate, since micronutrient antioxidants provide defense against tissue injury, high dietary intake of these micronutrients could be helpful to protect against osteoarthritis.

5. LOCAL RISK FACTORS

5.1. Joint injury and trauma

Articular cartilage tolerates loading from daily physical activities, in joints injuries and trauma the cartilage loses its flexibility, kills the cells and decrease the loading of the subchondral bone.²²

5.2. Obesity

People with an elevated body mass index (BMI) as a measure of relative weight for obesity, has a positive association between obesity and knee OA results in substantial overloading and damage to the knee joint.²³

5.3. Occupation

The lifting of heavy loads was found mainly in farmers, fishermen, construction site workers, and general laborers. Walking up stairs was experienced mainly by general laborers; all of these stress activities causes the strong association between knee injury and osteoarthritis.²⁴

5.4. Physical activity/Sports

Men & women practicing gymnastic or kung fu (martial arts) regularly were at the risk of Knee injury.²⁵

Schematic diagram of risk factors in osteoarthritis is shown in 2.




Fig. 2. Schematic diagram of risk factors for osteoarthritis.




Fig. 3. Potential targets for development of osteoarthritis in knee joint.

Osteoarthritis, or degenerative joint disease, is the most common form of arthritis. It mainly affects middle-aged and older people, involving the neck, lower back, knees, hips & fingers. Treatment aimed at control of pain using NSAIDs and physiotherapy. Conventional medicines do not prevent progression of osteoarthritis.

COMPOSITION OF ARTHRAID TABLET:



Each film coated tablet contains:



Supplement Facts	
Each film coated tablet contains (approx):	
Rose-Hip extract	250 mg
Collagen Peptide Type I	40 mg
Sodium Hyaluronate	30 mg
Chondroitin Sulfate Sodium	200 mg
Vitamin C	35 mg

INDICATIONS:

Osteoarthritis, rheumatoids arthritis, tendonitis and bursitis, muscular soreness and athletic injuries, carpal tunnel syndrome post-traumatic inflammation and pain.

CONTRA-INDICATIONS:

Known contraindications to any ingredients of the supplement.

Arthraid tablet might interact with drug thinning agents such as coumarins & warfarin.

DOSAGE AND DIRECTIONS FOR USE:

Take 1 tablet 1-2 times daily for up to three months. This intake may have to be adjusted for obese individuals including those taking diuretics, thereafter one tablet per day as a nutritional maintenance, as the arthritic condition improves. It is taken preferably with meals or as directed by a physician, licensed nutritionist. or certified trainer

PHARMACOLOGY:

ROSE HIP EXTRACT

- Rose hip extract is a herbal medicine from the Rosaceae family which contains polyphenols and anthocyanins.
- helps in preventing joint inflammation and damage and break down of cartilage.

COLLAGEN PEPTIDE

- Collagen peptide is a hydrolyzed form of collagen , a fibrous protein present in the extracellular matrix of living cells. Collagen peptide contains all essential amino acids except tryptophan.
- Helps in stimulating cartilage matrix there by regenerating joint cartilage.

SODIUMN HYALURAONATE

- It is a sodium salt of hyaluronic acid , glycosaminoglycan found in various connective, epithelial and neural tissues.
- It acts like a lubricant, helps in leading and stimulating growth of chondrocyte. Helps to treat osteo arthritis patients.

CHONDROITIN SULPHATE SODIUM

- It is one of the natural glycosaminoglycans, an important component of cellular matrix.
- It helps in production of hyaluronan which maintain viscosity in synovial fluid.
- Inhibits the enzymes leucocyte elastase and hyaluronidase.

VITAMIN C

- Acts both as antioxidant & metabolic activity.
- Helps in strengthen ligaments and tendons.
- Helps in neutralizes swelling and inflammation in osteoarthritis.

SAFETY:

ARTHRAID TABLET has an excellent safety record in both animal & human investigations, should be considered as a supplement of choice for nutritional correction of rheumatic disorders, can be safely taken alongside orthodox pain controlling medications.

ARTHRAID TABLET is generally regarded as safe when taken in the recommended doses; however, mild reactions can include gastrointestinal problems, such as nausea.

ARTHRAID TABLET is generally well tolerated. Because of lack of long-term safety data, ARTHRAID TABLET should be avoided by pregnant women and nursing mothers.

SIDE-EFFECTS:

Epigastric pain/tenderness, heartburn, diarrhea and nausea, flushing.

SPECIAL PRECAUTIONS:

Take ARTHRAID TABLET with or directly after meals to lessen the possibility of gastrointestinal upset. It should be avoided by pregnant women and nursing mothers.

STORAGE CONDITIONS:

Store in a cool & dry place, protected from light.

Keep out of reach of children.

STORAGE LIFE IS 2 YEARS.

The preparation should not be used after the expiry date.

6. CONCLUSION

Treatment of osteoarthritis (OA) is mainly based on the pathophysiological events that alter the initiation and progression of OA. Understanding the mechanism and Modulation of cytokines and MMPs would be a main target for treatment and prevention of Osteoarthritis. Conventional medicines do not prevent progression of osteoarthritis. Natural supplements like ARTHRAID TABLET have gained importance as a better alternative for the prevention of osteoarthritis progression.

REFERENCES

- 1.D.D. Dunlop, *et al.* Impact of joint impairment on disability-specific domains at four years; *Journal of Clinical Epidemiology*, 51 (12) (1998), pp. 1253-1261.
- 2.M. Steultjens, *et al.* Range of joint motion and disability in patients with osteoarthritis of the knee or hip; *Rheumatology*, 39 (9) (2000), pp. 955-96.
- 3.K.A. Sluka Pain mechanisms involved in musculoskeletal disorders; *Journal of Orthopaedic and Sports Physical Therapy*, 24 (1996), pp. 240-254.
- 4.N.B. Lawand, T. McNearney, K.N. Westlund Amino acid release into the knee joint: key role in nociception and inflammation; *Pain*, 86 (1) (2000), pp. 69-74
- 5.L. Arendt-Nielsen, *et al.* Sensitization in patients with painful knee osteoarthritis; *Pain*, 149 (3) (2010), pp. 573-581
- 6.S.R. McFaull, M. Lamontagne *In vivo* measurement of the passive viscoelastic properties of the human knee joint; *Human Movement Science*, 17 (2) (1998), pp. 139-165.

- 7.R.J. Johns, V. Wright Relative importance of various tissues in joint stiffness; *Journal of Applied Physiology*, 17 (5) (1962), pp. 824-828.
- 8.B. Hills, K. Thomas Joint stiffness and 'articular gelling': inhibition of the fusion of articular surfaces by surfactant; *Rheumatology*, 37 (5) (1998), pp. 532-538.
9. Schwarz, B. Hills Surface-active phospholipid as the lubricating component of lubricin; *Rheumatology*, 37 (1) (1998), pp. 21-2.
- 10.N. Smidt, *et al.* Effectiveness of exercise therapy: a best-evidence summary of systematic reviews; *Australian Journal of Physiotherapy*, 51 (2005), pp. 71-83.
- 11.N.A. Segal, *et al.* Quadriceps weakness predicts risk for knee joint space narrowing in women in the MOST cohort; *Osteoarthritis and Cartilage*, 18 (6) (2010), pp. 769-775
- 12.A. Young Current issues in arthrogenous inhibition; *Annals of the Rheumatic Diseases*, 52 (11) (1993), pp. 829-834
- 13.D.A. Rice, P.J. McNair, G.N. Lewis; *Arthritis Res Ther*, 1 (5) (2011), p.14
14. Qi-ping, *et al.* Clinical observation on treatment of 60 cases of osteoarthritis of knee joint by electroacupuncture; *Journal of Acupuncture and Tuina Science*, 1 (4) (2003), pp. 38-40
- 15.N. Verzijl, *et al.* *Arthritis & Rheumatism*, 46 (1) (2002), pp. 114-123
- 16.F.J. Keefe, *et al.* The relationship of gender to pain, pain behavior, and disability in osteoarthritis patients: the role of catastrophizing Pain, 87 (3) (2000), pp. 325-334
- 17.F. Cicuttini, T. Spector Osteoarthritis in the aged. Epidemiological issues and optimal management; *Drugs & Aging*, 6 (5) (1995), p. 409.
- 18.T.D. Spector, A.J. Mac Gregor Risk factors for osteoarthritis: genetics; *Osteoarthritis and Cartilage*, 12 (2004), pp. 39-44
- 19.D. Hunter, *et al.* Genetic contribution to bone metabolism, calcium excretion, and vitamin D and parathyroid hormone regulation; *Journal of Bone and Mineral Research*, 16 (2) (2001), pp. 371-378
- 20.R. Hirsch, *et al.* Osteoarthritis: new insights; *Ann Intern Med*, 133 (2000), pp. 635-646
- 21.C. Nishida, *et al.* The joint WHO/FAO expert consultation on diet, nutrition and the prevention of chronic diseases: process, product and policy implications; *Public Health Nutrition*, 7 (2004), pp. 245-250
- 22.J.A. Buckwalter, T.D. Brown Joint injury, repair, and remodeling: roles in post-traumatic osteoarthritis; *Clinical Orthopaedics and Related Research*, 423 (2004), pp. 7-16
- 23.A.T. Toivanen, *et al.* Obesity, physically demanding work and traumatic knee injury are major risk factors for knee osteoarthritis—a population-based study with a follow-up of 22 years; *Rheumatology*, 49 (2) (2010), pp. 308-314
- 24.P. Manninen, *et al.* Physical workload and the risk of severe knee osteoarthritis; *Scandinavian Journal of Work, Environment & Health* (2002), pp. 25-32
- 25.B. Järholm, *et al.* Age, bodyweight, smoking habits and the risk of severe osteoarthritis in the hip and knee in me; *European Journal of Epidemiology*, 20 (6) (2005), pp. 537-542